

METEOROLOGISK INSTITUTT
Norwegian Meteorological Institute

Transboundary air pollution by sulphur, nitrogen, ozone and particulate matter in 2020

The European Union

H. Klein, M. Gauss, S. Tsyro, Á. Nyíri, D. Heinesen, and H. Fagerli

Data Note 2022

ISSN 1890-0003

Contents

1	Introduction	3
1.1	The chapters of this report	3
1.2	Web interface for comparison with observations	4
1.3	Country codes	5
1.4	Definitions, statistics used	6
2	Emissions	8
2.1	Emissions used in the EMEP MSC-W model calculations	8
3	Time series	9
4	Transboundary fluxes	11
4.1	Deposition of oxidised sulphur	11
4.2	Deposition of oxidised nitrogen	12
4.3	Deposition of reduced nitrogen	13
5	Transboundary concentrations of ozone	14
5.1	MM-AOT40f	14
5.2	POD _{1.0,gen-DF} – Ozone fluxes to deciduous forests	15
5.3	SOMO35 – Risk of ozone damages to human health	16
6	Transboundary concentrations of particulate matter	17
7	Comparison with observations	19
8	Risk of damage from ozone and particulate matter in the European Union	23
8.1	Ecosystem-specific AOT40 values	23
8.2	Ecosystem-specific ozone fluxes	23
8.3	Health impacts from ozone and particulate matter	23

1 Introduction

This report is one in a series of country-specific notes, complementary to the EMEP Status Report 1/2022. It presents an overview of transboundary pollution of sulphur, nitrogen, ozone and particulate matter for the European Union in 2020.

All model runs for 2020, as well as the trend runs, have been performed with the EMEP MSC-W model version rv4.45, using ECMWF-IFS meteorology. The transboundary contributions presented here are based on source-receptor calculations with the EMEP MSC-W model for 2020, using meteorological and emission data for the year 2020.

As a basis for their correct interpretation, this section briefly explains what types of results are shown in this report and how they have been calculated.

1.1 The chapters of this report

Emissions (*Chapter 2*): The emissions for 1990–2020 have been derived from the 2022 official data submissions to UNECE CLRTAP as of 1 June 2022. The spatial (gridded) distribution of the emissions has been provided by the EMEP Centre on Emission Inventories and Projections (CEIP).

The gridded emission data used in the model calculations are available on WebDab at: <https://www.ceip.at/webdab-emission-database/emissions-as-used-in-emep-models>

For the years 2005–2020, emissions of condensable organics from residential combustion have been included based on the EMEPwRef2_v2.1C dataset from TNO (Netherlands Organisation for Applied Scientific Research). For more details please consult Appendix *Trend simulation done in 2022* in the EMEP Status Report 1/2022.

Time series (*Chapter 3*): Time series in depositions and air concentrations are presented for the period of 1990–2020. For all years, the meteorology of the respective year is used. Thus, interannual variability in the model results is due to changes in both emissions and meteorology. Please note that the time series are displayed for *EU-27* for the entire time period. Changes in membership during the period are not accounted for in these plots. Furthermore, the emission data and model version are updated regularly (see respective chapters on emissions and model updates in the EMEP Status Report 1/2022), which may lead to differences between results reported here and in earlier reports.

Transboundary fluxes (*Chapter 4*): Data are presented in the form of maps and pie charts. The data are generated by source-receptor calculations, where emissions for each emitter of one or more precursors are reduced by 15%. The results have been scaled up to represent the entire emission from an emitter.

Transboundary concentrations (*Chapters 5 and 6*): Data are presented in the form of maps and bar charts. Ozone and particulate matter are subject to significant non-linearities in chemistry. Therefore we calculate the effect of 15% reductions in emissions only.

The horizontal maps show the reduction in concentrations when emissions are reduced by 15% in the European Union. By convention, reductions in concentrations are represented by positive values in the maps. Thus, any negative values mean that concentrations increase as a result of an emission reduction (due to non-linearities in chemistry).

The bar charts identify the six most important emitter countries in terms of their effects on concentrations in the European Union that would result from a 15% reduction in emissions. In the bar charts, the sum of the *absolute values* of these effects corresponds to 100%. The percentage values (vertical scale in the bar charts) thus give an indication of the relative

importance of the various emitter countries that influence concentrations in the European Union (positive or negative, large or small contributions). Again, reductions are represented by positive values. Hence, a negative bar in the chart means that a *reduction* in emissions from an emitter country would lead to an *increase* in concentration in the European Union. In some countries this can occur because of strong non-linearities in chemistry.

In addition, for PM_{2.5} and PM₁₀ we show the total concentrations along with the percentage contribution from natural sources (sea salt and natural dust) to the total concentrations.

In the figures for ozone, we do not show contributions from areas that are outside the EMEP domain. Until 2019 these had been included as BIC (Boundary and Initial Conditions) and were calculated by reducing NO_x and NMVOC at the model boundary. However, the most important contributor to ozone from areas outside the EMEP domain is ozone itself, transported hemispherically across the model boundary. Including the BIC contribution that is due to NO_x and NMVOC only would be misleading. In principle, the BIC contribution due to hemispherically transported ozone could be included, but we choose here to focus on the contribution from countries within the EMEP domain.

Comparison with observations from the EMEP network (*Chapter 7*): The map of monitoring stations shows stations of the European Union in the EMEP measurement network with measurements in 2020 submitted to EMEP. The frequency analysis plots compare daily observation results with the model results. The measurement data are available from CCC:

<http://ebas.nilu.no>

The table provides annual statistics of the comparison of model results with observations for each measured component. Comparison is done only for stations with a sufficiently consistent set of data available in monthly or higher time resolution.

Risks from ozone and PM (*Chapter 8*): Particularly relevant for health impact, model results for SOMO35 (an ozone indicator) and particulate matter concentrations are shown. However, the results correspond to regional background levels and are not representative of local point measurements where these values can be much higher (e.g. in cities).

1.2 Web interface for comparison with observations

A more detailed evaluation against measurements from both the EMEP network and the European Environment Agency's (EEA) Air Quality e-Reporting Database can be found at the AeroVal web interface that has been developed recently:

<https://aeroval.met.no/evaluation.php?project=emep>

On that page the user can select the set of measurement data, the station or country of interest, and view a large number of statistical parameters (bias, correlation, root mean square error, etc.).

The web interface displays co-located observational and model datasets and contains:

- daily and monthly time series for each station, averaged per country, or the whole area covered by the model and the measurement network;
- statistics and scatter plots calculated for each station and country;
- an overall evaluation of the results using statistics calculated for each country or the whole area covered by the model and the measurement network (so-called Heatmaps and Taylor Diagrams).

Evaluation is made for O₃, PM_{2.5}, PM₁₀, SO₂, SO₄, NO₂, and several other nitrogen-containing species. The different types of visualization (bar charts, line charts, tables, etc.) are available both for viewing and for download.

1.3 Country codes

Many tables and graphs in this report make use of codes to denote countries and regions in the EMEP area. Table 1 provides an overview of these codes and lists the countries and regions included in the source-receptor calculations for 2020.

Code	Country/Region/Source	Code	Country/Region/Source
AL	Albania	IS	Iceland
AM	Armenia	IT	Italy
AST	Asian areas	KG	Kyrgyzstan
AT	Austria	KZ	Kazakhstan
ATL	N.-E. Atlantic Ocean	LI	Liechtenstein
AZ	Azerbaijan	LT	Lithuania
BA	Bosnia and Herzegovina	LU	Luxembourg
BAS	Baltic Sea	LV	Latvia
BE	Belgium	MC	Monaco
BG	Bulgaria	MD	Moldova
BIC	Boundary/Initial Conditions	ME	Montenegro
BLS	Black Sea	MED	Mediterranean Sea
BY	Belarus	MK	North Macedonia
CH	Switzerland	MT	Malta
CY	Cyprus	NL	Netherlands
CZ	Czechia	NO	Norway
DE	Germany	NOA	North Africa
DK	Denmark	NOS	North Sea
DMS	Dimethyl sulfate (marine)	PL	Poland
EE	Estonia	PT	Portugal
ES	Spain	RO	Romania
EU	European Union (EU27)	RS	Serbia
EXC	EMEP land areas	RU	Russian Federation
FI	Finland	SE	Sweden
FR	France	SI	Slovenia
GB	United Kingdom	SK	Slovakia
GE	Georgia	TJ	Tajikistan
GL	Greenland	TM	Turkmenistan
GR	Greece	TR	Turkey
HR	Croatia	UA	Ukraine
HU	Hungary	UZ	Uzbekistan
IE	Ireland	VOL	Volcanic emissions

Table 1: Country/region codes used throughout this report.

1.4 Definitions, statistics used

The following definitions and acronyms are used throughout this note:

SOA - secondary organic aerosol, defined as the aerosol mass arising from the oxidation products of gas-phase organic species.

SIA - secondary inorganic aerosols, defined as the sum of sulphate (SO_4^{2-}), nitrate (NO_3^-) and ammonium (NH_4^+). In the EMEP MSC-W model SIA is calculated as the sum:
 $\text{SIA} = \text{SO}_4^{2-} + \text{NO}_3^- (\text{fine}) + \text{NO}_3^- (\text{coarse}) + \text{NH}_4^+$.

SS - sea salt.

MinDust - mineral dust.

PPM - primary particulate matter, originating directly from anthropogenic emissions. One usually distinguishes between fine primary particulate matter, $\text{PPM}_{2.5}$, with aerosol diameters below $2.5 \mu\text{m}$ and coarse primary particulate matter, $\text{PPM}_{\text{coarse}}$ with aerosol diameters between $2.5 \mu\text{m}$ and $10 \mu\text{m}$.

$\text{PM}_{2.5}$ - particulate matter with aerodynamic diameter up to $2.5 \mu\text{m}$. In the EMEP MSC-W model $\text{PM}_{2.5}$ is calculated as $\text{PM}_{2.5} = \text{SO}_4^{2-} + \text{NO}_3^- (\text{fine}) + \text{NH}_4^+ + \text{SS}_{2.5} + \text{MinDust}(\text{fine}) + \text{SOA}(\text{fine}) + \text{PPM}_{2.5} + 0.13 \text{NO}_3^- (\text{coarse}) + \text{PM}_{25\text{water}}$. ($\text{PM}_{25\text{water}} = \text{PM}$ associated water).

$\text{PM}_{\text{coarse}}$ - coarse particulate matter with aerodynamic diameter between $2.5 \mu\text{m}$ and $10 \mu\text{m}$. In the EMEP MSC-W model $\text{PM}_{\text{coarse}}$ is calculated as $\text{PM}_{\text{coarse}} = 0.87 \text{NO}_3^- (\text{coarse}) + \text{SS}(\text{coarse}) + \text{MinDust}(\text{coarse}) + \text{PPM}_{\text{coarse}}$.

PM_{10} - particulate matter with aerodynamic diameter up to $10 \mu\text{m}$. In the EMEP MSC-W model PM_{10} is calculated as $\text{PM}_{10} = \text{PM}_{2.5} + \text{PM}_{\text{coarse}}$.

SS_{10} - sea salt aerosol with diameter up to $10 \mu\text{m}$.

$\text{SS}_{2.5}$ - sea salt aerosol with diameter up to $2.5 \mu\text{m}$.

SOx - group of oxidized sulphur components (SO_2 , SO_4^{2-}).

NOx - group of oxidized nitrogen components (NO , NO_2 , NO_3^- , N_2O_5 , HNO_3 , etc.).

redN - group of reduced nitrogen components (NH_3 and NH_4^+).

SOMO35 is the Sum of Ozone Means Over 35 ppb, an indicator for health impact assessment recommended by WHO. It is defined as the yearly sum of the daily maximum of 8-hour running average over 35 ppb. For each day the maximum of the running 8-hours average for O_3 is selected and the values over 35 ppb are summed over the whole year. If we let A_8^d denote the maximum 8-hourly average ozone on day d , during a year with N_y days ($N_y = 365$ or 366), then SOMO35 can be defined as:

$$\text{SOMO35} = \sum_{d=1}^{d=N_y} \max(A_8^d - 35 \text{ ppb}, 0.0)$$

where the \max function evaluates $\max(A - B, 0)$ to $A - B$ for $A > B$, or zero if $A \leq B$, ensuring that only A_8^d values exceeding 35 ppb are included. The corresponding unit is ppb.days.

AOT40 is the accumulated amount of ozone over the threshold value of 40 ppb, i.e.:

$$\text{AOT40} = \int \max(O_3 - 40 \text{ ppb}, 0.0) dt$$

where the \max function ensures that only ozone values exceeding 40 ppb are included. The integral is taken over time, namely the relevant growing season for the vegetation concerned, and in some daytime period. The corresponding unit is ppb.hours (abbreviated to ppb.h). The usage and definitions of AOT40 have changed over the years though, and also differ between UNECE and the EU.

Although the EMEP model generates a number of AOT-related outputs, we will concentrate in this report only on two definitions:

MM-AOT40f - AOT40 calculated for forests using estimates of O_3 at forest-top (*uc*: upper-canopy). This AOT40 is that defined for forests by the UNECE Mapping Manual, but using a default growing season of April-September.

MM-AOT40c - AOT40 calculated for agricultural crops using estimates of O_3 at the top of the crop. This AOT40 is close to that defined for agricultural crops by the UNECE Mapping Manual, but using a default growing season of May-July, and a default crop-height of 1 m.

POD_Y - Phyto-toxic ozone dose, is the accumulated stomatal ozone flux over a threshold Y , i.e.:

$$\text{POD}_Y = \int \max(F_{st} - Y, 0) dt \quad (1)$$

where stomatal flux F_{st} , and threshold, Y , are in $\text{nmol m}^{-2} \text{s}^{-1}$. This integral is evaluated over time, from the start of the growing season (SGS), to the end (EGS).

For the generic crop and forest species, the suffix *gen* can be applied, e.g. $\text{POD}_{Y,gen}$ is used for forests. POD was introduced in 2009 as an easier and more descriptive term for the accumulated ozone flux.

2 Emissions

2.1 Emissions used in the EMEP MSC-W model calculations




Figure 1: Spatial distribution of emissions from the European Union in 2020.

3 Time series

Important: For correct interpretation of the results shown in this chapter please read the paragraphs on *Emissions* and *Time series* in Section 1.1.

	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020
SO _x	21438	14311	8949	7084	3671	2461	2093	2054	1904	1655	1452
NO _x	15027	13031	11458	10660	8510	7140	6923	6785	6530	6173	5497
NH ₃	4841	4015	3975	3765	3557	3561	3577	3593	3552	3472	3441
NMVOC	16040	12785	10650	9080	7668	6644	6595	6633	6469	6360	6247
CO	56691	45249	34020	27913	24523	19562	19278	19142	18461	17619	16180
PM _{2.5}	3039	2327	1884	1843	1784	1500	1472	1456	1406	1318	1275
PM ₁₀	4881	3371	2728	2658	2496	2151	2116	2093	2058	1960	1905

Table 2: Emissions from the European Union. Unit: Gg. (SO_x given as SO₂, and NO_x as NO₂).

	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020
SO _x dep.	6990	4773	3194	2399	1632	1115	1031	984	979	909	743
NO _x dep.	3042	2766	2507	2302	1986	1620	1619	1592	1559	1455	1317
redN dep.	2752	2382	2353	2202	2184	2079	2157	2153	2125	2035	2012

Table 3: Estimated deposition of Sulphur (S) and Nitrogen (N) in the European Union. Unit: Gg(S) or Gg(N).

	1990	1995	2000	2005	2010	2015	2016	2017	2018	2019	2020
mean ozone	33	34	33	34	33	34	33	34	34	34	33
max ozone	43	43	42	42	41	42	41	42	42	42	41
MM-AOT40f	28017	25434	23171	21959	19338	18109	16122	16703	18997	17902	15863
SOMO35	3200	2902	2792	2737	2411	2424	2194	2348	2555	2429	2256
POD _{1.0,gen-DF}	31	30	29	28	27	24	25	24	27	19	19
PM _{2.5} anthrop.	13	11	9	9	7	7	6	6	6	6	6
PM ₁₀ anthrop.	17	13	12	11	9	9	8	8	8	8	8

Table 4: Estimated yearly mean values of air quality indicators averaged over the European Union. Unit: daily mean ozone (ppb), daily max ozone (ppb), MM-AOT40f (ppb·h), SOMO35 (ppb·d), POD_{1.0,gen-DF} (mmol/m²), PM_{2.5} (μg/m³) and PM₁₀ (μg/m³).




Figure 2: Annual emissions of photo-oxidant pollution precursors. Unit: Gg (note that NO_x is here given as NO₂).




Figure 3: Annual emissions and depositions of oxidised sulphur, oxidised nitrogen and reduced nitrogen. Unit: Gg(S) or Gg(N).




Figure 4: Changes in ozone related pollution relative to 2000. Unit: %. The large changes from year to year in some countries are mainly related to meteorological variability.




Figure 5: Mean concentrations of particulate matter. Unit: $\mu\text{g}/\text{m}^3$.

4 Transboundary fluxes

4.1 Deposition of oxidised sulphur




Figure 6: Contribution of emissions from the European Union to deposition of oxidised sulphur in the EMEP domain. Unit: $\text{mg(S)}/\text{m}^2$. The pie chart shows the six main receptor areas where oxidised sulphur from the European Union is deposited. Unit: %.




Figure 7: Top left: Deposition of oxidised sulphur in the European Union. Unit: $\text{mg(S)}/\text{m}^2$. Top right: The six main contributors to oxidised sulphur deposition in the European Union. Unit: (%). Bottom left: Oxidised sulphur deposition from transboundary sources. Unit: $\text{mg(S)}/\text{m}^2$. Bottom right: Fraction of transboundary contribution to total deposition. Unit: %.

4.2 Deposition of oxidised nitrogen




Figure 8: Contribution of emissions from the European Union to deposition of oxidised nitrogen in the EMEP domain. Unit: mg(N)/m^2 . The pie chart shows the six main receptor areas where oxidised nitrogen from the European Union is deposited. Unit: %.




Figure 9: Top left: Deposition of oxidised nitrogen in the European Union. Unit: mg(N)/m^2 . Top right: The six main contributors to oxidised nitrogen deposition in the European Union. Unit: %. Bottom left: Oxidised nitrogen deposition from transboundary sources. Unit: mg(N)/m^2 . Bottom right: Fraction of transboundary contribution to total deposition. Unit: %.

4.3 Deposition of reduced nitrogen




Figure 10: Contribution of emissions from the European Union to deposition of reduced nitrogen in the EMEP domain. Unit: mg(N)/m^2 . The pie chart shows the six main receptor areas where reduced nitrogen from the European Union is deposited. Unit: %.




Figure 11: Top left: Deposition of reduced nitrogen in the European Union. Unit: mg(N)/m^2 . Top right: The six main contributors to deposition of reduced nitrogen in the European Union. Unit: %. Bottom left: Deposition of reduced nitrogen from transboundary sources. Unit: mg(N)/m^2 . Bottom right: Fraction of transboundary contribution to total deposition. Unit: %.

5 Transboundary concentrations of ozone

5.1 MM-AOT40f




Figure 12: Reduction in MM-AOT40f that would result from a 15% reduction in emissions of NO_x (left) and NMVOC (right) from the European Union. Unit: ppb.h.




Figure 13: The six most important emitter countries or regions, with respect to their effects on MM-AOT40f in the European Union that would result from reductions in NO_x emissions (left) and NMVOC emissions (right).

5.2 $POD_{1.0,gen-DF}$ – Ozone fluxes to deciduous forests




Figure 14: Reduction in $POD_{1.0,gen-DF}$ that would result from a 15% reduction in emissions of NO_x (left) and NMVOC (right) from the European Union. Unit: mmol/m².




Figure 15: The six most important emitter countries or regions, with respect to their effects on $POD_{1.0,gen-DF}$ in the European Union that would result from reductions in emissions of NO_x (left) and NMVOC (right).

5.3 SOMO35 – Risk of ozone damages to human health




Figure 16: Reduction in SOMO35 that would result from a 15% reduction in emissions of NO_x (left) and NMVOC (right) from the European Union. Unit: ppb·day.




Figure 17: The six most important emitter countries or regions, with respect to their effects on SOMO35 in the European Union that would result from reductions in emissions of NO_x (left) and NMVOC (right).

6 Transboundary concentrations of particulate matter




Figure 18: Reduction in concentrations of SIA (left) and $PPM_{2.5}$ (right) that would result from a 15% reduction in emissions from the European Union. Unit: $\mu\text{g}/\text{m}^3$. Note the difference in colorbars.




Figure 19: The six most important emitter countries or regions, with respect to their effects on SIA (left) and $PPM_{2.5}$ (right) in the European Union that would result from reductions in emissions.




Figure 20: Left: PM_{10} concentration, and right: fraction of natural contributions of PM_{10} (sea salt and natural dust) to total PM_{10} concentration in the European Union. Units: $\mu\text{g}/\text{m}^3$ (left), % (right).




Figure 21: Left: Reduction in concentrations of $PM_{2.5}$ (left) and PM_{coarse} (right) that would result from a 15% reduction of emissions from the European Union. Unit: $\mu g/m^3$. Note the difference in colorbars.




Figure 22: The six most important emitter countries or regions, with respect to their effects on $PM_{2.5}$ (left) and PM_{coarse} (right) in the European Union that would result from reduction in emissions.




Figure 23: Left: $PM_{2.5}$ concentration, and right: fraction of natural contributions of $PM_{2.5}$ (sea salt and natural dust) to total $PM_{2.5}$ concentration in the European Union. Units: $\mu g/m^3$ (left), % (right).

7 Comparison with observations




Figure 24: Location of stations in the European Union.




Figure 25: Frequency analysis of ozone in the European Union at the stations that reported O₃ for 2020 (Observations, Model).




Figure 26: Frequency analysis of depositions in precipitation in the European Union (Observations, Model).




Figure 27: Frequency analysis of air concentrations in the European Union (Observations, Model).

Component	No.	Bias%	Correlation	RMSE
SO2 in Air	49	-23%±45%	0.38±0.34	0.15±0.10
Sulfate in Air	40	-39%±22%	0.67±0.30	0.59±0.38
NO2 in Air	58	-7%±40%	0.61±0.31	0.54±0.51
NO3- in Air	13	12%±53%	0.51±0.37	0.05±0.03
NH3+NH4+ in Air	28	-21%±35%	0.56±0.31	0.44±0.38
PM10	52	-29%±14%	0.54±0.29	4.93±2.35
PM2.5	43	-15%±21%	0.58±0.31	2.65±1.68
Ozone daily max	93	3%±5%	0.93±0.05	3.46±1.11
Ozone daily mean	93	6%±12%	0.91±0.07	4.48±1.93
SO4 wet dep.	48	-16%±27%	0.51±0.29	0.22±0.14
Nitrate wet dep.	48	21%±42%	0.47±0.28	0.44±0.25
Ammonium wet dep.	48	28%±46%	0.59±0.23	0.25±0.12
Precipitation	61	12%±21%	0.75±0.24	0.93±0.50

Table 5: Annual statistics of comparison of model results with observations in the European Union for stations with a sufficiently consistent set of data available in monthly or higher time-resolution. Standard deviations provide variability ranges between stations.

8 Risk of damage from ozone and particulate matter in the European Union

8.1 Ecosystem-specific AOT40 values




Figure 28: MM-AOT40f and MM-AOT40c in the European Union in 2020. *MM-AOT40f*: growing season April-September, critical level for forest damage = 5000 ppb·h; *MM-AOT40c*: growing season May-July, critical level for agricultural crops = 3000 ppb·h.)

8.2 Ecosystem-specific ozone fluxes




Figure 29: $POD_{3.0,gen-CR}$ and $POD_{1.0,gen-DF}$ in the European Union in 2020. Unit: mmol/m².

8.3 Health impacts from ozone and particulate matter




Figure 30: Left: Regional scale SOMO35, and right: PM_{2.5} in the European Union in 2020. SOMO35 is given in ppb·h, while PM_{2.5} concentrations are given in µg/m³.